

Table 3G
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Racial and Ethnic Distribution by Metro Area

Page: 1
Run Date: 7 / 2 / 12

Comparison of Racial and Ethnic Distribution (2)								
Metro Area	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Dane Cty	2,053	41.8	21.9	24.7	0.6	8.3	2.6	0.0
Mil. Cty	9,577	17.3	49.7	25.8	0.4	4.6	2.1	0.0
Racine	1,378	39.6	27.0	29.8	0.0	1.0	2.7	0.0
Rest of WI	20,013	72.4	3.6	13.8	2.5	5.2	2.4	0.0
Unknown Metro Areas	88	*	*	*	*	*	*	*
Wisconsin	33,109	53.3	19.0	18.6	1.7	5.0	2.4	0.0
Nation (Prior Year)	1,162,961	41.6	22.7	29.2	1.1	3.4	1.3	0.7

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4G
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Age Distribution by Metro Area

Page: 2
Run Date: 7 / 2 / 12

Comparison of Age Distribution (2)								
Metro Area	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Dane Cty	2,053	0.3	3.6	9.2	58.2	27.1	1.7	0.0
Mil. Cty	9,577	0.4	5.9	11.5	60.0	20.7	1.4	0.0
Racine	1,378	0.4	5.2	13.4	59.3	20.3	1.4	0.0
Rest of WI	20,013	0.1	3.8	11.5	63.5	19.7	1.3	0.0
Unknown Metro Areas	88	*	*	*	*	*	*	*
Wisconsin	33,109	0.2	4.4	11.4	62.0	20.5	1.4	0.0
Nation (Prior Year)	1,162,961	0.3	5.3	11.5	59.4	21.7	1.7	0.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5G
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Maternal Health Indicators by Metro Area

Page: 3
Run Date: 7 / 2 / 12

Comparison of Maternal Health Indicators (2, 3)															
Metro Area	Prepregnancy BMI (4)				Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Dane Cty	2,027	2.9 (1)	55.6 (2)	1,482	21.4 (1)	50.0 (2)	364	23.1 (1)	1,616	17.2 (1)	1,772	6.7 (4)	1,772	4.6(1)	
Mil. Cty	9,486	3.3 (2)	59.6 (3)	8,112	23.4 (4)	48.7 (1)	1,393	42.0 (4)	8,050	31.4 (4)	8,735	5.6 (2)	8,735	5.8(3)	
Racine	1,367	3.7 (3)	60.1 (4)	1,220	21.8 (3)	51.6 (4)	180	35.6 (3)	1,093	25.6 (3)	1,248	5.6 (2)	1,248	6.8(4)	
Rest of WI	19,769	3.8 (4)	55.5 (1)	15,234	21.7 (2)	50.0 (2)	2,921	26.1 (2)	15,815	18.9 (2)	18,096	5.3 (1)	18,096	4.6(1)	
Unknown Metro Areas	86	* *	* *	85	* *	* *	12	* *	75	* *	83	* *	83	* *	
Wisconsin	32,735	3.6	56.9	26,133	22.2	49.7	4,870	30.8	26,649	22.9	29,934	5.5	29,934	5.0	
Nation (Prior Year)	1,106,370	4.5	53.4	835,927	21.5	48.0	164,201	33.9	346,981	28.1	419,956	5.2	467,675	6.3	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6G
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Maternal Behavioral Indicators by Metro Area

Page: 4
Run Date: 7 / 2 / 12

Comparison of Maternal Behavioral Indicators												(2, 3)				
Metro Area	Medical Care						WIC Enrollment			Multivitamin Consumption						
	1st Trimester			No Care			1st Trimester			Pregpregnancy (4)			During Pregonancy (5)			
	Number	% (Rank)	% (Rank)				Number	% (Rank)		Number	% (Rank)		Number	% (Rank)		
Dane Cty	1,758	91.5	(1)	0.8	(1)		2,046	32.3	(4)		1,857	18.0	(2)	1,648	87.1	(2)
Mil. Cty	8,920	85.8	(3)	2.4	(3)		9,554	35.8	(2)		9,427	12.9	(4)	7,522	83.9	(3)
Racine	1,213	84.4	(4)	1.5	(2)		1,374	35.2	(3)		1,289	17.1	(3)	1,078	82.1	(4)
Rest of WI	18,428	89.5	(2)	2.4	(3)		19,952	36.5	(1)		19,383	21.5	(1)	15,823	87.4	(1)
Unknown Metro Areas	87	*	*	*	*		88	*	*		85	*	*	61	*	*
Wisconsin	30,406	88.3		2.3			33,014	36.0			32,041	18.6		26,132	86.2	
Nation (Prior Year)	429,770	83.1		4.1			1,074,863	36.0			380,154	19.2		418,448	85.0	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7G
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Smoking Indicators by Metro Area

Comparison of Smoking Indicators (2, 3)										
Metro Area	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)		Number	% (Rank)	Number	% (Rank)	
Dane Cty	1,983	28.2 (2)	1,765	14.6 (3)	376	35.9 (3)	1,674	11.5 (1)	1,767	2.9 (1)
Mil. Cty	8,936	22.5 (1)	8,731	11.3 (1)	1,497	38.1 (2)	7,591	16.3 (3)	8,734	6.4 (3)
Racine	1,320	33.3 (3)	1,244	14.3 (2)	312	40.1 (1)	1,081	19.0 (4)	1,245	6.5 (4)
Rest of WI	19,043	40.5 (4)	18,064	22.4 (4)	5,979	29.8 (4)	16,068	16.2 (2)	18,070	4.3 (2)
Unknown Metro Areas	84	* *	83	* *	28	* *	61	* *	83	* *
Wisconsin	31,366	34.3	29,887	18.3	8,192	32.1	26,475	16.1	29,899	4.9
Nation (Prior Year)	814,631	23.0	715,034	12.1	137,883	38.1	660,570	16.3	562,126	15.6

- (1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 8G
2011 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Infant Health Indicators by Metro Area

Page: 6
Run Date: 7 / 2 / 12

Comparison of Infant Health Indicators (2, 3)											
Metro Area	Birthweight					Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)								
	Number	% (Rank)	% (Rank)			Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Dane Cty	1,709	8.1 (2)	9.9 (4)			1,773	8.6 (1)	1,563	3.6 (3)	1,742	82.1 (1)
Mil. Cty	8,464	10.1 (4)	5.7 (1)			8,734	10.1 (3)	7,625	4.3 (4)	8,573	67.6 (4)
Racine	1,199	8.1 (2)	6.1 (2)			1,246	10.5 (4)	1,080	2.9 (2)	1,212	69.6 (3)
Rest of WI	17,518	6.8 (1)	8.1 (3)			18,157	8.9 (2)	15,938	2.6 (1)	17,865	73.7 (2)
Unknown Metro Areas	83	*	*	*	*	85	*	*	71	*	*
Wisconsin	28,973	7.9	7.4			29,995	9.3	26,277	3.2	29,477	72.3
Nation (Prior Year)	880,933	8.0	6.7			948,286	10.7	749,107	3.4	871,811	68.2

- (1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3G
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Racial and Ethnic Distribution by Metro Area
Combined 3 Years

Page: 7
Run Date: 7 / 2 / 12

Comparison of Racial and Ethnic Distribution (2)								
Metro Area	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Dane Cty	6,247	41.3	22.2	25.4	0.6	7.9	2.6	0.0
Mil. Cty	30,142	17.5	49.6	26.0	0.6	4.5	1.8	0.0
Racine	4,271	40.6	26.5	29.5	0.1	0.6	2.6	0.0
Rest of WI	62,046	72.9	3.4	13.9	2.6	4.9	2.2	0.0
Unknown Metro Areas	88	*	*	*	*	*	*	*
Wisconsin	102,794	53.5	19.1	18.8	1.8	4.7	2.1	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4G
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Age Distribution by Metro Area
Combined 3 Years

Page: 8
Run Date: 7 / 2 / 12

Comparison of Age Distribution (2)								
Metro Area	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Dane Cty	6,247	0.3	4.2	10.2	59.3	24.3	1.6	0.0
Mil. Cty	30,142	0.4	6.8	12.5	59.2	19.8	1.3	0.0
Racine	4,271	0.5	6.0	12.6	60.9	18.6	1.2	0.0
Rest of WI	62,046	0.1	4.2	12.4	63.3	18.7	1.2	0.0
Unknown Metro Areas	88	*	*	*	*	*	*	*
Wisconsin	102,794	0.3	5.0	12.3	61.7	19.4	1.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5G
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Maternal Health Indicators by Metro Area
Combined 3 Years

Page: 9
Run Date: 7 / 2 / 12

Comparison of Maternal Health Indicators (2, 3)															
Metro Area	Prepregnancy BMI (4)				Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Dane Cty	6,159	3.1 (1)	54.7 (1)	4,432	21.4 (2)	50.7 (3)	1,141	25.2 (1)	4,818	20.2 (2)	5,516	6.3 (4)	5,516	5.2(2)	
Mil. Cty	29,838	3.2 (2)	58.8 (3)	24,191	24.2 (4)	48.1 (1)	4,654	43.1 (4)	25,182	32.6 (4)	27,531	4.9 (1)	27,531	6.5(3)	
Racine	4,230	3.3 (3)	58.9 (4)	3,810	21.5 (3)	51.8 (4)	627	34.1 (3)	3,412	25.0 (3)	3,874	5.3 (3)	3,874	6.7(4)	
Rest of WI	61,319	3.8 (4)	54.8 (2)	46,825	21.3 (1)	50.6 (2)	9,442	26.6 (2)	48,729	19.8 (1)	56,315	5.0 (2)	56,315	5.0(1)	
Unknown Metro Areas	86	* *	* *	85	* *	* *	12	* *	75	* *	83	* *	83	* *	
Wisconsin	101,632	3.6	56.2	79,343	22.2	49.9	15,876	31.6	82,216	24.0	93,319	5.1	93,319	5.5	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6G
2009 - 2011 Pregnancy Nutrition Surveillance
Wisconsin
Comparison of Maternal Behavioral Indicators by Metro Area
Combined 3 Years

(1)

Page: 10
Run Date: 7 / 2 / 12

Comparison of Maternal Behavioral Indicators												(2, 3)				
Metro Area	Medical Care						WIC Enrollment			Multivitamin Consumption						
	1st Trimester			No Care			1st Trimester			Pregpregnancy (4)			During Pregarancy (5)			
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	
Dane Cty	5,542	90.9	(1)	1.0	(1)		6,228	31.1	(4)		5,656	17.8	(2)	4,915	86.3	(2)
Mil. Cty	28,128	83.4	(4)	2.8	(4)		30,035	35.1	(3)		29,584	12.4	(4)	23,718	82.4	(4)
Racine	3,843	83.7	(3)	2.5	(2)		4,261	36.8	(1)		4,121	16.6	(3)	3,391	83.8	(3)
Rest of WI	57,244	88.6	(2)	2.5	(2)		61,851	35.7	(2)		60,041	21.1	(1)	48,719	86.5	(1)
Unknown Metro Areas	87	*	*	*	*		88	*	*		85	*	*	61	*	*
Wisconsin	94,844	87.0		2.5			102,463	35.3			99,487	18.1		80,804	85.2	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7G
2009 - 2011 Pregnancy Nutrition Surveillance
Wisconsin
Comparison of Smoking Indicators by Metro Area
Combined 3 Years

(1)

Page: 11
Run Date: 7 / 2 / 12

Comparison of Smoking Indicators (2, 3)										
Metro Area	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Dane Cty	6,020	29.5 (2)	5,506	15.2 (2)	1,237	39.2 (1)	4,992	13.2 (1)	5,508	3.5 (1)
Mil. Cty	28,120	23.1 (1)	27,520	11.7 (1)	4,932	38.3 (2)	23,947	18.8 (3)	27,527	7.5 (3)
Racine	4,099	32.4 (3)	3,870	15.2 (2)	981	35.7 (3)	3,424	20.8 (4)	3,871	8.3 (4)
Rest of WI	59,187	41.3 (4)	56,208	22.9 (4)	18,799	29.8 (4)	49,391	17.9 (2)	56,240	4.9 (2)
Unknown Metro Areas	84	* *	83	* *	28	* *	61	* *	83	* *
Wisconsin	97,510	35.0	93,187	18.8	25,977	32.1	81,815	18.0	93,229	5.7

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 8G
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Infant Health Indicators by Metro Area
Combined 3 Years

Page: 12
Run Date: 7 / 2 / 12

Comparison of Infant Health Indicators (2, 3)											
Metro Area	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)		High (5)								
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Dane Cty	5,346	7.5 (2)	9.0 (4)		5,550	8.7 (1)	4,882	2.9 (3)	5,443	81.6 (1)	
Mil. Cty	26,593	9.8 (4)	5.9 (1)		27,526	10.8 (4)	23,745	4.2 (4)	26,956	66.0 (3)	
Racine	3,721	8.1 (3)	6.0 (2)		3,869	10.1 (3)	3,352	2.8 (2)	3,766	64.8 (4)	
Rest of WI	54,453	6.9 (1)	8.1 (3)		56,409	9.1 (2)	49,422	2.6 (1)	55,367	72.0 (2)	
Unknown Metro Areas	83	*	*	*	*	85	*	*	71	*	*
Wisconsin	90,196	7.9	7.4		93,439	9.6	81,472	3.1	91,617	70.5	

- (1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

